








J. W. DAWSON.

PICTOU, N. S.





Pictou Literary and Scientific Society.

"With the starting of the Grammar School in Pictou, and the foundation of the Academy, the nineteenth century saw the beginning of the Pictou County renaissance; and we find in 1834 the foundation of the Pictou and Literary Society.

... The program included a weekly lecture by a member. J. D. B. Fraser and John W. Dawson, in the early years, contributed most valuable scientific papers.

The Society continued to prosper for many years. It did incalculable good in Pictou and Pictou County. It gave Sir J. W. Dawson his early opportunity to speak of his reading and early geological discoveries."

From "The Pictou Advocate,"

Pictou, Nova Scotia,

28 December, 1933.

App M 105^A/₃₄

Essays

(on)

Geology.

Read before the Pictou Literary Society
March 1836

Those parts of the Earth beneath its surface, to which mankind have been able to penetrate, have in all ages been objects to which their attention has been particularly directed, both from a curiosity of knowing the structure of the globe, and from the utility of the minerals which they contained. It was not, however, till the 14 century, that any attempt was made to reduce the knowledge of these to scientific principles. At that period the attention of the learned in Europe, was directed to this subject, by the discovery of shells and bones imbedded in the rocks at the foot of the

the Asperines; and many curious opinions were entertained respecting their origin. One class of disputants maintained that these shells and bones never had belonged to animals, but were formed spontaneously in the rock in which they were found, by what they called "plastic force"; or a tendency in stones and clays to arrange themselves in certain definite forms. Others asserted that all those rocks which contain animal or vegetable remains, such as limestone, sandstone, and coal, were the residuum of the deluge. And others that they were all formed by the action of fire. These, and many other opinions just as remote from truth, were discussed for more than a century; and then, after much opposition, the principles which form the basis of

of modern systems of geology were established. These are that the nature and

These principles the
however are re- mal
and used,
not comprehension and the
open many opinions have
been adopted as s, tr.
feeling them that ing
in a short essay le-
such as that which the
essay I have to offer only
a general outline of things
to present ourselves
appears to give a general map

of rocks, clay, sand, and other heterogeneous materials, heaped together without order or regularity. When however we examine the crust of the globe, in those places where it has been laid open by mines and other excavations, we

we discover that it is composed of a great number of layers of rock varying in thickness from a few inches to many hundred feet and each of which is by some peculiarity distinguished from those immediately adjoining it. These have been termed the strata of the earth. Imbedded in the rocks belonging to these strata there is an immense variety of remains of animals and vegetables which have existed in former ages of the world and belonging to species the far greater part of which have now no living representatives.

They thus afford in characters which can never be obliterated a record of the animals and vegetables which have existed upon earth to a period far antecedent to the era of history.

These fossils are called Organic Remains. And it is the investigation of the origin nature and
 cor-

arrangement of the strata and the organic remains contained in them which is the province of Geology.

This science has often been confounded with mineralogy it will therefore be necessary to state the distinction between them. Mineralogy refers to the chemical composition and more minute peculiarities of rocks and earths and Geology to the general structure of the earth and to the positions in it occupied by the strata. This subject may therefore be ~~considered~~ divided into two parts 1st Those features which belong to all the strata of which the earth is composed and 2^{dly} the order in which they occur in the crust of the globe and the peculiarities which distinguish them from each other.

If we dig into the earth after passing through the vegetable soil

we find sand and clay containing loose blocks of stone; and below these hard rocks in layers parallel to each other. These are composed of very few materials the principal of which are Quartz or Flint Clay and Limestone but these are mingled in such a manner as to produce a considerable variety of combinations. They are also arranged in a certain determinate order which is never in any degree inverted. Thus if we suppose the strata to be represented by the letters of the Alphabet A being that nearest the surface and Z the lowest; rocks of the same kind as A are never found below Z or any of the intervening letters. The knowledge of this principle is of great advantage in mining thus if a person searching for coal discover upon the surface any rock

rock known to be below it in the series he is certain that no coal exists there. The reverse of this however is no sure ^{ground} of information as it often happens that in particular places many steps in the series are wanting this is caused by the circumstances in which all the strata were originally formed, that is ~~that~~ the materials of which they are composed were washed by water from the land which existed at the period of their formation and deposited at the bottom of seas oceans and lakes.

It would thus happen that those parts which were nearest the land would consist of much coarser materials than that which had been in the ~~middle~~ ^{centre}. That all the ^{another ~~series~~ is the different elevations} stratified rocks were deposited ^{and depressed to which the strata have} by the action of water is one of the ^{best suggest} fundamental principles of Geology. and it is corroborated

by the appearances of the strata themselves as for example their containing pebbles having the appearance of having been long tossed about in the sea, immense quantities of shells and corals which from their perfect state and the positions in which they are found ~~has~~ must have lived in the same places in which they are discovered and ~~from~~ logs of wood and blocks of stone with marine shell fish adhering to them

From this it is evident that they must have been deposited upon a foundation which was level or nearly so But at present they are found not level or in the bottom of the sea but upon dry land and inclined at different angles twisted and contorted and some times even in a vertical position and
it

it is to this disarrangement that we owe the greater part of the hills mountains and valleys with which the surface of the earth is diversified.

The question thus arises how were these rocks elevated from the bottom of the ocean and from the level state to the positions which they now occupy? and Geologists from a careful examination of the strata themselves and of the effects at present produced by volcanic agency have referred it to this cause. The correctness of this conclusion is proved by the existence of what are called

Unstratified Rocks.

These are not placed in layers like the others but are generally found in masses in columns or in upright walls called dikes. They contain no pebbles or organic remains, are very hard and many of them in² mineralogical structure

bear a striking resemblance to the
 lavas of modern volcanoes. The most
 common of these are granite whin-
stone and basalt. They are al-
 ways found below the stratified
 except where they have protruded
 through them from beneath. Their posi-
 tions and some of the disturbances
 to which strata are subject will
^{see page} be seen by a diagram. In No 1
 A B & C represent unstratified rocks
 which have forced themselves through
~~the~~ strata lying above them A & B have
 sent out shoots or veins into the stra-
 tified rock and A & C after reaching
 the top have flowed over No 2 repre-
 sents strata which have been twist-
 ed by force from below while in a soft
 state. In No 3 the strata marked
 A have been driven through that
 marked B while that marked
 I has been formed after the dis-
 tur-

bance took place. Though the diagram is imaginary cases nearly resembling it are often found in nature as will be seen from a section of part of the Coast of England. In No I the unstratified rocks under A have placed the strata B in a ~~horizontal~~^{vertical} position, and in No II the strata c and unstratified rock d have forced the coal bed E in to a curve and G are the remains of strata which have been formed after the disturbance took place but have been subsequently washed away.

From such facts as these it is evident that the unstratified rocks were forcibly ejected from the earth in a melted state. This is rendered still more certain by the effects which they have produced in the strata through which they have passed. Thus where they have passed

sect

sed through coal they have converted it into a substance resembling coke and where they have passed through chalk it has become as hard as marble

Since therefore these unstratified rocks have been thus geeted and in almost every case where elevation of strata has been observable it has been produced by them we may fairly conclude that all the strata were elevated by the same cause. It is however proved by the appearance of the strata themselves that they were elevated from the sea in small portions and at far distant periods This accounts for some of the oldest strata appearing on the surface ~~while~~ at one place while at another not far distant ^{and} ~~far more recent~~ strata may occur. As for example

some

of strata belonging to the oldest strata being found at the head of the East and Middle Rivers while at the Albion mines the coal which is far more recent occurs. Having thus given a short sketch of the general features of the strata the next point is the Order in which they occur in the crust of the Earth and the peculiarities by which they are distinguished. In classifying the strata Geologists found that the lowest or most ancient division of rocks with which they were acquainted contained no pebbles or organic remains. This therefore they called the Primary Strata. Above this was another class which did contain organic remains but these belonged to species of animals which are now extinct. This they called the Secondary strata. Another class yet

more recent was also discovered containing some remains of animals and plants of the same species which now exist. This they called the Tertiary strata. And the accumulation of sand clay and loose stone immediately below the vegetable soil was called Deluvium. Each of these divisions I shall therefore consider in their order beginning with the

Primary Strata.

These rocks contain no organic remains or pebbles. This however does not prove that there never existed in them as as many of the rocks composing them appear to have been melted and all of them have been subjected to immense pressure. They consist principally of limestone flint slate gneiss and other hard rocks ~~and~~ they contain all the metals except mercury and many of

the finest beds of *Stetucary Marble*.
 The granite rocks on the southern
 coast of Nova Scotia belong to this
 class. The next general division
 is

The Secondary Strata

This contains a great variety of rocks
 formed from those parts of the more an-
 cient strata which at the period of
 their formation were elevated above the
 level of the sea and exposed to the
 action of the elements. They have been
 subdivided into a number of groups each
 of which contains many beds of rock.

The first and lowest of these is the
Greywacke Group.— In this occur the first
 signs of animal life in the shape of
 a small species of shell-fish called *Tri-*
lobites and a few corals. No remains of
 any terrestrial animals have been found
 in this and some of the strata above it
 and from this circumstance many
 geologists have concluded that at the
 pe-

riod when they were formed the whole earth was covered with water. This however is unfounded as we know that at present notwithstanding the numbers of animals in existence the bones of very few are deposited in the ~~body~~ of the ocean and as all the strata were formed in the bottom of the sea the same will apply to them.

To this group the clay slate upon which the town of Halifax is built and which extends along the head of the East and Middle rivers and occurs in Antigonish is supposed to belong. ^{and in it are the remains of iron ore, of the east river,} Above this is the Old Red Sandstone. This is principally composed of the coarse red sandstone from which it derives its name and which ^{derives} ~~derives~~ its colour from the oxide of Iron. It contains many beds of water-worn pebbles but few organic remains.

next above this is the
 Mountain Limestone. — This rock is very
 plentiful it contains veins of spar and
 in many places is wholly composed
 of shells and corals. ~~It~~ ^{They} contains va-
 luable mines of lead It underlies the
 Pictou coal-field. The next group is the
 Coal Measures. — This is one of the most
 important of the strata. it does not
 consist entirely of coal but is inter-
 stratified with many beds of sand-
 stone and shale often containing
 iron. The coal itself ~~is~~ ^{mostly} is entirely
 composed of vegetable ^{matter} which does
 not however as has sometimes been
 supposed grow in the ground but
 is the remains of forests which have
 grown upon the earth in former ages
 but from the pressure to which it
 has been subjected it has been reduced
 to an undistinguishable mass in which
 however traces of leaves and other remains
 yet exist

and which when chemically analysed is found to consist of the same materials as wood. Coal beds are at present forming in many places particularly at the mouths of great rivers where the vegetable matter washed down at once time and the sand and mud at another may produce the same alterations which are observed in the ancient coal measures.

One of the most curious facts with respect to the coal strata is that at the period of their formation the climates of Europe and America even in their most northern latitudes were as warm as those of tropical countries are at present This is proved by the palm trees, tree ferns and other productions of warm latitudes which are found in them and which from their perfect state would

not

have been carried from any great distance. The coal also bears evidence of having been formed in straits between small islands and thus when ~~they~~^{it} was formed the greater part of the continents of Europe and America consisted of archipelagos similar to those of the Pacific ocean.

In ^{to the local} local beds are often found "faults" or breaks in the strata ^{which} ~~called~~ in which it often happens that one part of the strata has been elevated above or depressed below the other. These faults frequently occur in the Pictou coal-field. The next is the

Red Marl group. - This is often of great thickness. It is composed of beds of different kinds of sandstone limestone and mud-dy sediment and sometimes ^{contains} rock salt. The red sandstone of river John belongs to this group and the salt springs at the head of the West River are

also supposed to belong to it. It is therefore probable that extensive beds of rock salt may exist at the salt springs. Among the organic remains found in this group are those of two gigantic species of marine animals called the

Ichthyosaurus and *Plesiosaurus*

The former of these is about 32 feet ^{in length} with a long snout and 60 teeth in each jaw and 4 fins resembling those of a whale. Like this animal it seems to have been formed for living in the waters of the ocean and to have been able to swim with great velocity.

The latter is remarkable for the length of its neck which exceeds that of any other known animal having no less than 41 joints. It seems to have been adapted for living in shallow water

such

as the margins of rivers and lakes
 Both the *Tethiosaurus* and *Plesiosaurus*
 from the undigested bones
 found within their skeletons seem
 to have preyed upon fish and
 occasionally to have devoured
 the young of their own species. ^(digested)

Above the Red Marl is the
 Colite Group. — This contains a great
 variety of sandstones and claystones
 and a kind of limestone in oval
 blocks resembling eggs from which
 the group derives its name. There
 has been much disputing as to what
 could provide the material for this
 limestone; some have referred it
 to shells and corals and others to
 mineral sponges. In this group
 are found the *Tethiosaurus* and
Plesiosaurus and remains of
 a great number of curious reptiles,
 a few vegetables of kinds very
 dif-

ferent from those now existing
and great numbers of shellfish
Of these latter one kind called
the Ammonite of Cornu Am-
monis is so abundant that beds
of considerable thickness are formed
of the shells of this species alone
Above this is the

Chalk Group. - The greater part
of this group consists of the well-
known mineral from which it de-
rives its name although in its low-
er parts there are extensive deposits
of sandstone sand and clay. In
its upper parts the chalk is mix-
ed with flints and in its lower parts
these are wanting. The chalk
belonging to the group has there-
fore been divided into Chalk with
flints and chalk without flints.
This group appears to be entire-
ly wanting in Nova Scotia. It

Corn

contains many organic remains. The vegetables ~~which~~ are principally marine but some terrestrial reptiles are also found. Having thus shortly noticed the groups which compose the secondary strata we must next consider the third great division or the

Tertiary Strata.

The existence of these strata was discovered by Currier who found that the city of Paris was built upon a kind of basin surrounded by chalk hills which had been gradually filled up with thick beds of sandstone clay and limestone more recent than ^{any} the secondary strata and containing a few bones of animals belonging to species the same as now exist. He therefore divided them into a separate group. Currier's discovery having directed the attention of other Geologists to these strata it was afterwards
found

that the valley of the Thames was
 a basin similar to that of Paris and
 subsequently that tertiary strata were
 found in different parts of Europe
 These strata are not like the sec-
 ondary deposited continuously de-
 posited over large tracts of Country
 but are found in basins of com-
 paratively small extent which seem
 at some period to have formed the
 beds of small seas and inland seas
^{in which the tertiary strata were formed and}
 through this is the most recent
^{at some times to have been covered with}
 of those rocks which may properly
^{salt and at other not put up.}
 be termed stratified there is a group
 above it which as it contains an im-
 mense quantity of animal remains
 some of which do not differ in many
 respects from those now inhabiting
 the earth is worthy of notice. This
 is the accumulation of sand clay
 gravel and pebbles which lies im-
 mediately below the vegetable soil

It appears to consist of fragments which have been conveyed to their present position by water and from this many geologists have concluded that it is the residuum of the deluge. The organic remains which it contains are principally those of terrestrial animals; of these the principal is the Mammoth which seems to have been a large species of Elephant distinguished from those which at present exist by the superior curvature of its tusks and by its being covered with hair. A perfect specimen of this animal was found enclosed in an iceberg (where it had probably remained since the deluge) at the mouth of the river Lena in Siberia. Its flesh was so well preserved that dogs and wolves collected from all quarters to feed upon it. A rather gigantic animal.

which is found in this formation
 is the *Megatherium* an animal
 about 17 feet long and 7 in height
 It appears to have nearly resembled
 the ant-eater or sloth and to have
 been a burrowing animal. Along
 with the bones of this animal were
 found parts of a large shell ~~which~~
 resembling that of the armadillo
 and which seems to have belong-
 ed to the *Megatherium*. Many con-
 jectures have been made as to what
 advantage an animal so large
 could have derived from a shell.
 Some have supposed that its
 use was to prevent the masses
 of earth which in burrowing might
 occasionally fall upon him from
 injuring his back; and others that
 it served as a protection from wild
 beasts. In this group the principal
 deposits of such bones are in caverns
 one of

of the principal of these is the Kirk-
 dale cave in ~~Yorkshire~~ In the bottom
 of which were found imbedded in hard
 mud the bones of the Elephant Rhino-
 ceros. Hippopotamus, Hyæna, Tiger, Bear,
 Horse, Wolf, Fox, Weasel, Ox, Deer, Hare,
 Rabbit, Water-rat, Mouse, Raven, Pigeon,
 Lark, and Thrush. All these bones were
 gnawed and broken and from this it is
 supposed that the cave was in ante-
 deluvian ages a den of hyænas and
 that the other animals were dragged in
 to it as food. It thus appears that ~~the~~
 Great Britain was at one time inha-
 bited by animals which ^{are} now found
 only in intertropical regions. There
 are besides the Cave of Kirkdale many
 other caverns ^{in Europe} containing bones in yet
 greater abundance. From one in Italy
 a ship's load was carried to France
 to be employed in refining sugar
 for which purpose however they
 were found to be useless. and in one of

the valleys at the foot of the
 et lps the leg bones and ribs of Eleph-
 ants were found in such abundance
 that the peasants used to employ
 them for building fences ~~This~~

I Have thus given a very brief ac-
 count of the strata in their order be-
 ginning with the oldest ~~X~~ before lea-
 ving them it must be remarked
 that the whole series never occurs
 in one place and that the very
 oldest are often seen on the surface
 of high mountains thus the High-
 lands of Scotland are almost whol-
 ly composed of the Primary Strata

From ^{diagram} what has been said res-
 pecting the structure of the Earth it
 will be seen that no bones either
 of Monkeys or of Men have been
 found in the strata. And from this
 it is evident that these animals
 either did not exist in the most
 ancient periods of the globe or

were confined to small limits. From this circumstance also and from the fact that in the more ancient strata only shell fish and other animals supposed to be of simple structure are discovered and as we reach the more modern groups we gradually find animals more complex in their structure. La March deduced the theory that at the creation only a few simple animals were formed and that these gradually improved themselves until a shellfish after passing through many gradations at length became a monkey an orang-outang and finally a man. An amusing refutation of this theory is given by Lyell who notices the custom of some of the natives of Sumatra who train monkeys to climb cocoa nut trees and throw down the nuts and then corks bore

these

these Indians after attempting in ^{vain}
 to climb the ^{rocks} ~~steep~~ ^{for their original shape} ~~and not acquire~~
 the long arms and ^{prehensile} feet with fingers
 and thumbs like the monkeys in-
 stead of being obliged to turn them
 inwards. It is also proved by
 the appearance of the strata that
 at the period of their formation the
 Northern parts of Europe and Am-
 erica enjoyed a climate as warm
 as that of tropical countries at the
 present day, ^{and that the heat gradually decreased} so account for this
 many theories have been advanced
 as for example that the poles of the
 Earth had changed their position
 and that the Earth was created
 red hot and had been gradually
 cooling. These however are utterly
 at variance with the common
 course of nature and therefore ex-
 tremely improbable. There is one
 opinion however first brought for-
 ward

would, by M^r Lyell which is very pro-
 bable and generally believed That the
 warmth of the Earth may at the
 time when the strata were formed
 may be accounted for by the dif-
 ferent positions of the Great continen-
 ts of the Globe. That this is capable
 of producing these effects may be
 seen from the comparative heat
 of Europe and America The former
 in corresponding latitudes is the war-
 mer and this because it is re-
 heated from the arctic regions by
 water while America stretches to
 the very pole ^{where} If then the great
 continents of the world were situated
 nearer the Equator than they are
 at present they themselves would
 be warmer than they ^{now} are at pre-
 sent and also the warmth of the
 whole globe would be increased
 another proof of this is that the
 coal-

coal-measures and others of the secondary strata of Europe and North America seem to have been formed among small Islands and therefore if there were then any great continents they must have been in different positions from those occupied by those now existing

Another and more interesting part of Geology is the consideration of the question whether the causes at present acting upon earth are capable of producing the stratified rocks which exist in the crust of the ^{globe} and what strata are at present forming. All stratified rocks must be formed from substances existing previous to their formation. Those causes therefore which produce strata must be such as either abstract matter from the interior of the

Earth and deposit it on its surface or ~~detach~~ wash earth and other substances from the land into the sea. These may be included under the heads of Springs, rivers, coral formations, tides and currents earthquakes and volcanoes.

1. Springs.— The causes of springs were fully explained in the lecture on Hydrostatics lately delivered before the society. It will therefore be necessary only to consider their effects in forming strata. It is a very rare occurrence to find a spring not in some degree impregnated with mineral matter which it is continually transferring from the crust of the earth to its surface some however contain it in very small quantities while others such as the West over salt springs contain very great proportions

All of these however Insignificant though their action may appear carry an immense quantity of matter to the sea or deposit it upon land. (£2,000 tons)

2. Rivers.— In dry weather rivers do not carry to the sea any very considerable deposits but in the spring and fall when the rivers are flooded and the ground at the same time rendered soft and brittle by the frost very great quantities of sand mud gravel and vegetable matter are conveyed to their mouths. These substances when deposited in water which is not frequently agitated generally accumulate in deposits called Deltas immediately at the mouths of rivers. Of the formation of these the Nile is a very good example It has not only deposited at its mouth a large

delta but has within the period of history elevated the part of Egypt which it overflows to the height of more than six feet

3 Coral reefs and Islands. The animals which form these through themselves very diminutive have continued on the earth longer than any other animals as their corals which are their workmanship are found on the very lowest of the secondary strata a great part of the limestones contained in which are to be attributed to their industry and they are yet found in great abundance and still continue to form great numbers of Islands and rocks. To these belong a great part of the Islands in the Pacific ocean Coral Islands generally appear above water in the form of a ring containing a small pool or lake of water

and having one or more openings. This circumstance together with the fact that coral insects never begin to build at any greater depth than from fifty to a hundred feet which proves that the coral islands must be built upon some foundation elevated above the bed of the surrounding sea. Induced Mr. Lyell to conclude that the circular form of coral islands was owing to their being built upon the margin of the craters of extinct submarine volcanoes. As coral insects never build above the level of the sea their islands never attain any great elevation unless raised by volcanic force. As soon however as they reach the level of the sea they become covered with a dense growth of vegetables and soon be-

come habitable. Some coral formations are of great extent: one upon the coast of Holland stretches to the length of 1000 miles

4 Tides and Currents. — It is a general principle in Geology that all land which is above the level of the sea is gradually wearing away and upon no part of its surface is this more apparent than upon its sea coasts. In such situations the decrease is caused by the waters of the sea which are continually opening the amount of clay land and depositing matter in in their own bed. ~~To this~~ Of this the East coast of Britain is an example. Its shores are wasted by the tides of the German ocean which have in the era of history very much changed the appearance of the coast. The sites of towns and villages have been

obliged to be moved farther inland
 producing valuable tracts of country
 have been destroyed.

5. Volcanoes and Earthquakes.— Although
 there are very few of what are com-
 monly called volcanoes volcanic force
 exerts far more power in the formation
 and arrangement of strata than might
 be supposed as it is of ten exerted with-
 out breaking out into eruptions
 A singular instance of this is the chain
 of the Andes and Rocky Mountains
 and indeed nearly all the great
 mountains of the world are of
 volcanic origin The whole num-
 ber of volcanoes upon earth is
 estimated at 200 of these Ameri-
 ca contains 116, Asia 66, and Eu-
 rope 18. None having yet been dis-
 covered in Africa. Very many
 of these are called extinct because
 no eruptions from them are known
 to have taken place in the period

of history from this however it does
 not follow that these volcanoes are
 incapable of causing eruptions for
 the periods of these in particular
 volcanoes are generally far distant
 and many which have been long
 considered extinct have given evi-
 dence of the existence of volcanic
 power in them. The mode in which
 volcanoes act is by streams of water
 and mud by lava or by showers of
 ashes and hot stones. The first of
 these which is rarest of the accom-
 panies the eruptions of particular
 volcanoes as of Cotopaxi in the An-
 des while in those of others it is en-
 tirely wanting. It is supposed to
 originate partly from water emit-
 ted from the earth and partly from
 the melting of snow on the summit
 of the mountains. In one eruption
 of the kind in Italy water was
 poured

from a rent in the side of a mountain, consuming numbers of small fish. The other substances ejected from volcanoes ~~are~~ lava and ashes are most frequent and destructive of man and his works and it is these which have given rise to the erroneous idea that volcanoes emit flame and smoke the former being the red gleam of the melted lava and hot stones and cinders and the latter clouds of steam or ashes.

~~The cones~~ Almost all volcanic mountains are formed of lava and ashes regularly stratified and intersected by lava dykes.

Volcanic action has been ascribed to various causes. Some have asserted that the interior of the earth was a mass of fire or of melted rocks and have adduced as a proof the fact that

as we descend into a mine the air becomes warmer. But it has been found that by allowing this heat to increase in the same proportion at a few miles below the surface we should find a temperature equal to the melting point of Iron and consequently should then arrive at the surface of the melted part of the earth which would thus be a greater proportion to the solid crust than the yolk of an egg to its shell and it is evident that such a crust could retain its solid state for a moment. It is therefore impossible that this can be the cause of volcanoes and besides it has never been explained how such heat could be produced.

That explanation which is generally believed is that volcanoes

oes are produced by chemical action, and from the circumstance that volcanoes are always found near the sea or in situations where they have access to water ~~the~~ is concluded that the combustion is produced by this agent. This is corroborated by the fact that there are many minerals ^{and compounds of them} which take fire by exposure to water.

Earthquakes are produced by the same causes as volcanoes and act upon the crust of the earth by heaving up particular parts and depressing others (clefts or faults).

It thus appears that the causes which have been mentioned are now forming strata similar to those which at present compose the crust of the

the earth containing remains
of Man and of his works which
may in future ages of the world
be viewed with the same cur-
iosity as that with which we
view the bones of the Mastodon
or the Mammoth

457

47.

The following pages are
evidently a Catalogue, describing
specimens collected by Sir J.
William Dawson in his youth.

The essay at the beginning of
this book was read in 1836 when
he was only 15 (his birthday being
in the autumn); and this collection
was probably begun about that
time, as each set of specimens
begins with number 1.

William B. Dawson,
(son of Sir J. W. D.)

May, 1934.

f-2

The following pages are evidently a Catalogue, describing specimens collected by Sir J. William Dawson in his youth.

The essay at the beginning of this book was read in 1836 when he was only 15 (his birthday being in the autumn); and this collection was probably begun about that time, as each set of specimens begins with number 1.

William B. Dawson,
(son of Sir J. W. D.)

May, 1934.

The following paper are
 evidently a Catalogue, describing
 specimens collected by Sir F.
 William Dawson in his youth.
 The essay at the beginning of
 this book was read in 1836 when
 he was only 12 (his birthday being
 in the autumn); and this collection
 was probably begun about that
 time, as each set of specimens
 begins with number 1.

William F. Dawson,
 (son of Sir F. W. D.)

Nov. 1834.

Multivalves

Genus 1 Chiton - valves placed in transverse pleats down the back. Shell in the form of a boat inverted and high at the stern.



2. Lepus - Barnacle - conical formed of triangular valves broad at the base and tapering at the top where they are united by horizontal plates.



3. *Pholus*. Stone Pierces Shell
bivalve, obliquely striated, with
several small accessory valves at
the hinge, an incurved tooth in-
side beneath the hinge.



2. *Tellina*. Hinging on one side
in the fore part of one valve
there is a convex in the other a
concave fold - 3 teeth lateral
ones smooth in one valve.

Shell often
delicate and
beautifully colored.

3. *Chelon*. Three species well
 known, especially *testudinatus*, with
 several small accessory values at
 the fringe, and numerous teeth in
 the side beneath the fringe.
 but seldom more than one or two
 inserted in the opposite row.

4. *Chelon*. Three species well
 known, especially *testudinatus*, with
 several small accessory values at
 the fringe, and numerous teeth in
 the side beneath the fringe.
 but seldom more than one or two
 inserted in the opposite row.

5. *Chelon*. Three species well
 known, especially *testudinatus*, with
 several small accessory values at
 the fringe, and numerous teeth in
 the side beneath the fringe.
 but seldom more than one or two
 inserted in the opposite row.

6. *Chelon*. Three species well
 known, especially *testudinatus*, with
 several small accessory values at
 the fringe, and numerous teeth in
 the side beneath the fringe.
 but seldom more than one or two
 inserted in the opposite row.

Bivalves

1 Mya - Gaper - Generally gaping at one end. Hinge with broad strong teeth seldom more than one & not inserted in the opposite valve.
brown coat



2 Tellina. Hinging on one side in the fore part of one valve there is a convex in the other a concave fold - 3 teeth lateral ones smooth in one valve.



Shells often delicate and beautifully colored

3. *Cardium*, *Coehle*. Equisilateral
 Equivalve, convex, ribbed striated
 or grooved longitudinally. toothed
 margin, two teeth near the beak
 and a larger one at each side.
 Shell thick often
 spherical




4. *Maestra* - Unequal sided equi-
 valve Middle tooth with a hol-
 low in each side lateral ones
 remote Shells usually thin & delicate




5. Donax. Wedge — bluntly toothed
margin — blunt or broad margin
Teeth / marginal one a little be
hind Sometimes smooth at
others shined & filed.



6. Solen. Razor fish — open at
both ends, often covered with a
thin cuticle



7 Verruc. frontal margin flattened. 3 teeth, lateral ones divergent at the tip. interior often coloured
Wampum—



S. Spondylus. Thorny oyster—beak solid with unequal valves one of the valves ~~flatter~~ the other nearly flat 2 teeth separated by a small hollow. Covered with longitudinal rows of spines.

7. *Persea*: fruit at an average flat
 toward 3 teeth. *Chloranthus* over *Chloranthus*
 cut the tip. *Chloranthus* after *Chloranthus*
Chloranthus —

11. *Asper*: *Chloranthus*. *Chloranthus* with
Chloranthus. *Chloranthus* after *Chloranthus* — *Chloranthus*
Chloranthus with *Chloranthus* *Chloranthus* over
 of the *Chloranthus* *Chloranthus* after *Chloranthus*
 flat. 2 teeth separated by a small
Chloranthus. Covered with *Chloranthus*
 room of *Chloranthus*.

9. Chama. Cum. Shell coarse
hinge with a hard gibbosity.
often not equilateral.

10. Arce. Equivalve. hinge with
numerous sharp teeth united be-
tween each other shell thick
sometimes covered with a cuticle
sometimes crenulated sometimes
grooved

8. Chamber. Glass. Shell coarse
 rings with a hard pithy
 often not separated.
 Chamber. Shell coarse

10. Acer. Cylindrical. Rings with
 numerous sharp teeth inserted
 between each other shell thick
 sometimes covered with a scale
 sometimes ornamented
 grooves around the rings
 of the scales distinct the other
 grooves distinct from the
 rings at some points

11 Ostrea — generally with unequal valves hinge without teeth but an ovate hollow transverse groove divided into Scallops and oysters

14 Perna — very brittle, large
hinged with no teeth — small

12 Anomia — not equi-valve
beak curved over the hinge one
of the valves flattish the other
gibbous. Shells thin sometimes
shaped at others smooth.



27. *Botrychium* generally with an
 equal number of stamens and
 pistils. The stamens are
 inserted into the ovary and
 the pistils are

28. *Botrychium* - The stamens are
 inserted into the ovary and
 the pistils are - but separate
 from the stamens over the large
 of the ovary. The stamens
 are inserted into the ovary and
 the pistils are

13 *Mytilus* *Muscle* — hinge usually without teeth, some species have grooves some are much others striated.

14 *Purra*. — Very brittle, large thickened ribs — no teeth — resembles a wing or fin.



23. *Mytilus edulis* — many
usually without teeth, some of
the large species have one or
two sharp teeth.

Mytilus — shell very
thin, with one or two
teeth, and a small
opening.

24. *Mytilus* — very little
known, with one or two
teeth — usually
one or two.

25. *Mytilus* — large
shell, with sharp
teeth.

Univalves

1. Shells having a spine

1 Argonauta. - Shell extremely thin, with only one compartment resembling a scroll with a large aperture.



2. Nautilus. Some spiral others tooth shaped.



3. Lower - Bone structure of skull
 different from that of the 1st.

1. Skull having a fissure

2. Asymmetrical. Skull extremely
 thin, with only one carpal bone
 resembling a scroll with a large
 aperture.
 4. Supraorbital - Curved, toothed.

2. Ventral - Bone of spine
 when tooth shaped.


3 Conus - Cone shaped smooth
without teeth. slit at the lip



4. Cypraea - Cury, toothed.




3. Caranx - Bone structure smooth
Dentition teeth. *but at the tip*
of the jaw



4. Polistes - Bone structure smooth
Dentition teeth. *but at the tip*
of the jaw



5. Buller - without teeth
 A pectine large at the base oblong



6 Voluta Shell one collected.
 teeth or pleats on the puller
 side of the lip.



2. *Strophomena* - small, thin, shell-like
 at the base of the columnar structure
 leading to the right with a small
 or irregular hollow top containing

3. *Strophomena* - shell are called
 teeth or points on the hollow
 side of the tip.

4. *Strophomena* - small, thin, shell-like
 at the base of the columnar structure
 leading to the right with a small
 or irregular hollow top containing

7 Buccinum - gibbous Aperture
 vate terminally in a short canal
 leading to the right with a leaf
 or projection full lip extended



8. Strombus - Aperture much
 dilated the lip expanding and
 produced into a groove.




The following is a list of the
 most important points in the
 history of the city of
 St. Petersburg, from its
 foundation to the present time.

8. The following is a list of the
 most important points in the
 history of the city of
 St. Petersburg, from its
 foundation to the present time.

9 Murex, - Trumpet shell
rough, Membranous, aperture
oval with a straight or slightly
ascending canal.



10. Trochus. Corneal, Aperture
angular or rounded upper side
contracted



11. Turbo. Wreath. Spiral long
Aperture contracted orbicular in
line



12. Helix. or Spiral - Delicate
little Aperture contracted - semicircular, or rounded



13. Nerita or Hoof-shell
 Gibbous flattish at bottom a-
 perture semilunar bilobed like
 truncate sometimes toothed
 aperture.



14. Heliotis or Ear shell -
 Ear shaped with a row of
 orifices along the surface hinge
 nearly concealed.



13. *Chloris a. chloris* (the same)
 14. *Chloris a. chloris* (the same)
 15. *Chloris a. chloris* (the same)
 16. *Chloris a. chloris* (the same)
 17. *Chloris a. chloris* (the same)


18. *Chloris a. chloris* (the same)
 19. *Chloris a. chloris* (the same)
 20. *Chloris a. chloris* (the same)
 21. *Chloris a. chloris* (the same)
 22. *Chloris a. chloris* (the same)

Without a Shine

15. *Petellae* or Limpet. Cone
basin shaped some have a pe-
tition and appear like a slipper
some are perforated at the apex



16. *Dentalium* or Tooth Shell
Tubular straight or slightly curved
with an undivided cavity open
at both ends.




17. *Tabellae* - Shell tubular com-
posed of several broken shells united
all substances & glued together.

17. *Portulaca oleracea* L.
Common in wet places, especially in the
lowlands and swamps. The leaves are
succulent and fleshy.

18. *Portulaca quadrifida* L.
Common in wet places, especially in the
lowlands and swamps. The leaves are
succulent and fleshy.


19. *Portulaca* ...
Common in wet places, especially in the
lowlands and swamps. The leaves are
succulent and fleshy.

17. Serpula or worm shell.
Tubular, adhering to other substances separated by ~~external~~ divisions sometimes spiral.




18 Terebra or Ship Worm. —

Shell tapering flexuous with two
Hemispherical ~~and~~ two cancellated
valves before



19. Sabella — shell tubular composed of sand broken shells, vegetable substances &c glued together



17. *Leptorhynchus* or worm shell
Leptorhynchus, exhibiting the other
 species separated by external
 characters.

18. *Leptorhynchus* or worm shell
 exhibiting the other
 species separated by external
 characters.

19. *Leptorhynchus* or worm shell
 exhibiting the other
 species separated by external
 characters.

- 1 Smooth Bivalves - Limestone from Cape Breton
- 2 Bivalves - striated longitudinally probably cardiac White Carb - Lime Pugwash
- 3 Bivalves deeply striated - ^{and Productus} cardiac - Black Carb Lime Pictou Island
- 4 Leaves & Stems of two species of Euerinites Do Do
- 5 Leaves of Ferns, Shale - New- road
- 6 Stems of Calamites Canadensis from Patterson Mills off Ken. Dies quarry and 6 mile brook

- 7 Stems of Smooth Succulent
vegetables shale & sandstone
New Road &c
- 8 Long lanceolate leaves
probably of Calamites Patter-
son Mills, Pictou, &c
- 9 Small Heart shaped leaves
Patterson Mills
- 10 Smooth lignite spotted like
Beech wood, sandstone, loc-
ality unknown.
- 11 ^a Three forked leaf like
that of Oxalis acetosella
* wood sorrel or Sheep grass.
but a little smaller. P.M.
* Club Pentagynia fern, diss., Geraniaceae

85-

12 Globules consisting of carb.
onaceous matter containing
oxide of Iron Pat-Mill
Sometimes sand or clay

13. Leaves of Grapes. P.M.

14 Parts of the impression of a
large plant 8 inches broad at
the battery there in a coarse
sandstone. The impression
was nearly a foot long and
in that distance there
was no perceptible nar-
rowing.

15 Pieces of a plant con-
verted into sulphured
and ochreous oxide of iron


found at Hutchens farm
 in a piece of shaly sandstone.
 It extended at right angles
 to the planes of stratification
 had a crooked root-like
 appearance, and no fresh
 white markings, the wood
 itself had a stringy ap-
 pearance.

16. A small ammonite
 apparently smooth about
 $1\frac{1}{4}$ inch in whole diameter
 and a little more than $\frac{1}{8}$ th
 of an inch in the diameter
 of the whorls, found in a
 black shaly piece of Pie-

97
Two Island Limestone - the
other in the same piece
- both destroyed

17. Irregular oblong pieces
of white lime or gypsum, about
2 inches long, and little more
than $\frac{1}{2}$ in. broad, $\frac{1}{8}$ thick
found in P & G lime, nature
unknown.

18. A large specimen of a
rough barked tree about
2 inches thick petrified into
some kind of calcareous
mass at the sides and by
into in the centre ~~petrified~~
the same as 15: found in

a hard ^{9.8.} grey conglomerate
~~like sandstone~~, at
mortimer's shore near Howlets
to the east of the same
place were strata
of sandstone containing

Calamites & smooth stems
with a stratum of hard grey
argillaceous conglomerate
like the same rocks at Bell

19. At Howlets shore in
the Sandstone before men-
tioned two smaller specimens
of No 14. In the grey con-
glomerate of same place
one calamite and small
pieces of coal or lignite, in

the sandstone smooth reeds
 like a specimen found at
 R. Is garden. No 7. Also, stem
 1 inch diameter striped trans-
 versely. ligneous impression
 In a hard blue sandstone
 a specimen of a large cactus,
 beautifully marked nearly
 2 inches diameter, surrounded
 by serbaceous matters. X

28 At Kitchen's farm. Leaves
 of a species of fern broad in-
 dented leaves.

X This appeared to be a
 ligneous impression, the
 cortical one was smooth

21 At Howlet's a stem

1 inch in diameter cortical
impression roughly striated
like a line thick, lyeuous
impression nearly smooth.

also a smaller specimen
of 19. in sandstone, blue

22 a stem shaped trunk
persey 1 inch diameter no
marks of leaves or branches

23. Two specimens of
a horn shaped plant
striped like calamites

2 inches thick much

flattened blue sandstone



at Houletts and detritus at
Battery,


Small round bodies resembling
leek bulbs to which Mr. P. M.

at Charlotte and detection at
 Battery, Charlotte, N.C.

found in dissected cortical
 fragments, mostly in the
 form of thin, thin, brown
 fragments, nearly smooth.
 also a smaller specimen
 of 19, in kind the same

22 a stem also at the
 base, each diameter no
 more than leaves or branches

23 Two specimens of
 a horn shaped plant
 tipped like calamities
 2 inches thick much
 flattened blue or indigo



104.

105-

106

107

108

109

100.

112

113

115

118.

118.

120

121

123

124

125-

126

127

128

138-

136

